

Sore Throat - Symptom

Definition

- Pain, discomfort or raw feeling of the throat
- Pain is made worse when swallows
- The infection is in the throat and usually also involves the tonsils.
- Children less than 2 years of age usually can't complain about a sore throat. A young child who does not want favorite foods may have a sore throat. They may also start to cry during feedings.

Health Information

Causes

- Usually caused by an infection of the throat (pharyngitis). Since the tonsils are in that area, they are usually also infected (tonsillitis).
- **Colds (URIs).** Most sore throats are part of a cold. In fact, a sore throat may be the only symptom for the first 24 hours.
- **Influenza.** A sore throat can be the first symptom in flu infections.
- **Viral pharyngitis.** Some viruses cause a sore throat without nasal symptoms.
- **Herpangina.** A special viral infection of the throat that causes little ulcers near the tonsils.
- **Strep pharyngitis.** Group A Strep is the most common bacterial cause. It accounts for 20% of persistent sore throats. Only these need an antibiotic.
- **Postnasal drip.** Drainage from sinus congestion can cause a sore throat. Sometimes, it's because the secretions are irritating to the throat. More often, it's from the frequent throat clearing.
- **Mouth breathing.** Children who sleep with the mouth open may wake with a sore throat. It feels better within an hour of having something to drink. Use a humidifier to help prevent a dry mouth.

Strep Throat

- Symptoms include sore throat, fever, headache, stomach pain, nausea and vomiting.
- Cough, hoarseness, red eyes, and runny nose are usually not seen with Strep throat. These symptoms point more to a viral cause.
- Scarlet fever rash (fine, red, sandpaper-like rash) is highly suggestive of Strep throat.
- Peak age: 5 to 15 years old. Not common under 2 years old unless sibling has Strep.
- Diagnosis should be confirmed by a Strep test before starting treatment. There is no risk to wait until a Strep test is done.
- Standard treatment is with antibiotics by mouth.

Prevention of Spread to Others

- Good hand washing can prevent spread of infection.

Care Advice

1. Overview:

- Most sore throats are just part of a cold and caused by a virus.
- A cough, hoarse voice or nasal discharge points to a cold as the cause
- If a sore throat is the only symptom, it's still usually viral. A fever or swollen neck lymph nodes point to this as the cause.
- Here is some care advice that should help.

2. **Sore Throat Pain Relief:**
 - Age over 1 year. Can sip warm fluids such as warm chicken broth or apple juice. Some children prefer cold foods such as popsicles or ice cream.
 - Age over 6 years. Can also suck on hard candy or lollipops. Butterscotch seems to help.
 - Age over 8 years. Can also gargle. Use warm water with a little table salt added. A liquid antacid can be added instead of salt. Use Mylanta or the store brand. No prescription is needed.
 - Medicated throat sprays or lozenges are generally not helpful.
3. **Pain Medicine:**
 - To help with the pain, give acetaminophen (such as Tylenol) or ibuprofen. Use as needed.
4. **Fever:**
 - For fevers above 102° F (39° C), give acetaminophen (such as Tylenol) or ibuprofen. Note: Lower fevers are important for fighting infections.
 - For ALL fevers: Keep your child well hydrated. Give lots of cold fluids.
5. **Antibiotics Not Needed:**
 - Antibiotics are not helpful for viral infections.
 - They can only kill bacteria.
6. **Fluids and Soft Diet:**
 - Try to get your child to drink adequate fluids.
 - Goal: Keep your child well hydrated.
 - Cold drinks, milk shakes, popsicles, slushes, and sherbet are good choices.
 - **Solids.** Offer a soft diet. Also avoid foods that need much chewing. Avoid citrus, salty, or spicy foods. Note: Fluid intake is much more important than eating any solids.
 - Swollen tonsils can make some solid foods hard to swallow. Cut food into smaller pieces.
7. **What to Expect:**
 - Most often, sore throats with a viral illness last 4 or 5 days.
8. **Return to School:**
 - Your child can return to school after the fever is gone.
 - Your child should feel well enough to join in normal activities.
 - Also, children with Strep throat need to be taking an antibiotic for 12 hours.

Call Your Doctor If

- Trouble breathing or drooling occurs
- Sore throat is the main symptom and lasts more than 48 hours
- Sore throat with a cold lasts more than 5 days
- Fever lasts more than 3 days
- You think your child needs to be seen
- Your child becomes worse

Pediatric Care Advice

Author: Barton Schmitt MD, FAAP

Copyright 2000-2023 Schmitt Pediatric Guidelines LLC

Disclaimer: This health information is for educational purposes only. You the reader assume full responsibility for how you choose to use it. The information contained in this handout should not be used as a substitute for the medical care and advice of your pediatrician. Listing of any resources does not imply an endorsement.