

IBUPROFEN (Medicine for pain or fever, such as Advil or Motrin)

DOSAGE TABLE

PRODUCT	CHILD'S WEIGHT (POUNDS)							
	12-17 lbs	18-23 lbs	24-35 lbs	36-47 lbs	48-59 lbs	60-71 lbs	72-95 lbs	96+ lbs
Infant's Liquid 50 mg / 1.25 mL	1.25 mL	1.875 mL	2.5 mL	3.75 mL				
Children's Liquid 100 mg / 5 mL			5 mL	7.5 mL	10 mL	12.5 mL	15 mL	20 mL
Children's Chewable 100 mg tablet			1 tab	1 ½ tabs	2 tabs	2 ½ tabs	3 tabs	4 tabs
Children's 100 mg tablet					2 tabs	2 tabs	3 tabs	4 tabs
Adult 200 mg tablet					1 tab	1 tab	1 tab	2 tabs

CHART NOTES

- **Brand names:** Advil, Motrin or store brand ibuprofen.
- **Dose:** Find your child's weight in the top row of the dosage table. Look below the correct weight for the dose based on the product you have. Adult dose is 400 mg.
- **Measure the Dose:** Use the syringe or dropper that comes with the medicine. If not, you can buy a medicine syringe at a drugstore. If you use a teaspoon, it must be a measuring spoon. **Reason:** Regular spoons are not reliable. Keep in mind: 1 level teaspoon equals 5 mL.
- **How Often:** Repeat every 6 to 8 hours as needed. Don't give more than 3 times a day.
- **Age Limit:** Don't use younger than 6 months unless told to by your child's doctor. **Reason:** For any fever in the first 12 weeks of life, your baby needs to be seen now. Also, the FDA has not approved ibuprofen for infants younger than 6 months.
- **Caution:** Do not use acetaminophen (Tylenol) and ibuprofen together. Reason: no benefit over using one medicine alone and a risk of giving too much. **Exception:** Your child's doctor told you to give both.

CALL YOUR DOCTOR BACK IF...

- Your child looks or acts very sick
- Any serious symptoms occur like trouble breathing
- Any fever occurs if under 12 weeks old
- Fever without other symptoms lasts over 24 hours (if age less than 2 years)
- Fever lasts over 3 days (72 hours)
- Fever goes above 104° F (40° C)
- You think your child is in pain
- You think your child needs to be seen
- Your child becomes worse

This handout is intended for informational purposes only and should not be used as a substitute for the care and advice of a medical professional. The accuracy of the information contained in this handout is not guaranteed and there may be variations in treatment that your doctor or nurse may recommend based on individual facts and circumstances.